

Via Aristotle, Leibnitz & Mach to a fractal $D = 2$ universe

D. F. Roscoe

3rd February 2008

Abstract

Over the past two decades or so, it has become increasingly apparent that, out to quite large distances, galaxies are distributed in a quasi-fractal fashion with fractal dimension $D \approx 2$. Whether or not this behaviour continues onto indefinitely large scales is a matter of live debate and is a question that can only be settled if, at some scale, there is an unambiguous transition to homogeneity. This point has not yet been reached and may never be. This paper has been written on the basis of the tentative hypothesis that quasi-fractal $D \approx 2$ behaviour is a persistent and fundamental feature of galaxy distribution on all scales and addresses the question of the origins of this putative fractality. Given this tentative hypothesis then, except for the device of putting the fractal behaviour into the initial conditions - which is to by-pass the question, there is no obvious explanation within the framework of conventional cosmology. So we adopt the position that *fractality* is per-se a signature that different thinking is required. We find that a beautiful solution flows deductively from elementary observations about the world in which we live when we take seriously a view of space and time that can be traced, via Mach, Berkeley and Leibnitz to Aristotle. In summary, we find that a globally inertial space and time can be irreducibly associated with a fractal, $D = 2$, distribution of material if we are prepared to review our understanding of what is meant by the notion of *metric* on astrophysical scales.

1 Introduction

A basic assumption of the *Standard Model* of modern cosmology is that, on some scale, the universe is homogeneous; however, in early responses to suspicions that the accruing data was more consistent with Charlier's conceptions of an hierarchical universe [1, 2, 3] than with the requirements of the *Standard Model*, de Vaucouleurs [4] showed that, within wide limits, the available data satisfied a mass distribution law $M \approx r^{1.3}$, whilst Peebles [5] found $M \approx r^{1.23}$. The situation, from the point of view of the *Standard Model*, has continued to deteriorate with the growth of the data-base to the point that, according to Baryshev et al [6]

...the scale of the largest inhomogeneities (discovered to date) is comparable with the extent of the surveys, so that the largest known structures are limited by the boundaries of the survey in which they are detected.

For example, several recent redshift surveys, such as those performed by Huchra et al [7], Giovanelli and Haynes [8], De Lapparent et al [9], Broadhurst et al [10], Da Costa et al [11] and Vettolani et al [12] etc have discovered massive structures such as sheets, filaments, superclusters and voids, and show that large structures are common features of the observable universe; the most significant conclusion to be drawn from all of these surveys is that the scale of the largest inhomogeneities observed is comparable with the spatial extent of the surveys themselves.

In recent years, several quantitative analyses of both pencil-beam and wide-angle surveys of galaxy distributions have been performed: three recent examples are give by Joyce, Montuori & Labini [13] who analysed the CfA2-South catalogue to find fractal behaviour with $D = 1.9 \pm 0.1$; Labini & Montuori [14] analysed the APM-Stromlo survey to find fractal behaviour with $D = 2.1 \pm 0.1$, whilst Labini, Montuori & Pietronero [15] analysed the Perseus-Pisces survey to find fractal behaviour with $D = 2.0 \pm 0.1$. There are many other papers of this nature in the literature all supporting the view that, out to medium depth at least, galaxy distributions appear to be fractal with $D \approx 2$.

This latter view is now widely accepted (for example, see Wu, Lahav & Rees [16]), and the open question has become whether or not there is a transition to homogeneity on some sufficiently large scale. For example, Scaramella et al [17] analyse the ESO Slice Project redshift survey, whilst Martinez et

al [18] analyse the Perseus-Pisces, the APM-Stromlo and the 1.2-Jy IRAS redshift surveys, with both groups finding evidence for a cross-over to homogeneity at large scales. In response, the Scaramella et al analysis has been criticized on various grounds by Joyce et al [19].

The argument has recently reduced to a question of statistics: basically, the proponents of the fractal view argue that the statistical tools (eg correlation function methods) widely used to analyse galaxy distributions by the proponents of the opposite view are deeply rooted in classical ideas of statistics and implicitly assume that the distributions from which samples are drawn are homogeneous in the first place. Thus, much effort is being expended developing tools appropriate to analysing samples drawn from more general classes of populations - a general focus being the idea that one should not discuss fractal structures in terms of the correlation amplitude since the only meaningful quantity is the exponent characterizing the fractal behaviour. Recent papers arguing this general point of view are Sylos Labini & Gabrielli [20] and Gabrielli & Sylos Labini [21].

This work of this paper is based upon the view that this putative fractal behaviour is fundamental and persistent on all scales, and is a signal that our conventional understanding of space & time needs to be reconsidered.

1.1 A brief history of ideas of space and time

The conception of space as the container of material objects is generally considered to have originated with Democritus and, for him, it provided the stage upon which material things play out their existence - *emptiness* exists and is that which is devoid of the attribute of *extendedness* (although, interestingly, this latter conception seems to contain elements of the opposite view upon which we shall comment later). For Newton [22], an extension of the Democritian conception was basic to his mechanics and, for him:

... absolute space, by its own nature and irrespective of anything external, always remains immovable and similar to itself.

Thus, the absolute space of Newton was, like that of Democritus, the stage upon which material things play out their existence - it had an *objective existence* for Newton and was primary to the order of things. In a similar way, time - *universal time*, an absolute time which is the same everywhere - was also considered to possess an objective existence, independently

of space and independently of all the things contained within space. The fusion of these two conceptions provided Newton with the reference system - *spatial coordinates* defined at a *particular time* - by means of which, as Newton saw it, all motions could be quantified in a way which was completely independent of the objects concerned. It is in this latter sense that the Newtonian conception seems to depart fundamentally from that of Democritus - if *emptiness* exists and is devoid of the attribute of *extendedness* then, in modern terms, the *emptiness* of Democritus can have no *metric* associated with it. But it is precisely Newton's belief in *absolute space & time* (with the implied virtual clocks and rods) that makes the Newtonian conception a direct antecedant of Minkowski spacetime - that is, of an empty space and time within which it is possible to have an internally consistent discussion of the notion of *metric*.

The contrary view is generally considered to have originated with Aristotle [23, 24] for whom there was no such thing as a *void* - there was only the *plenum* within which the concept of the *empty place* was meaningless and, in this, Aristotle and Leibnitz [25] were at one. It fell to Leibnitz, however, to take a crucial step beyond the Aristotelian conception: in the debate of Clarke-Leibnitz (1715~1716) [26] in which Clarke argued for Newton's conception, Leibnitz made three arguments of which the second was:

Motion and position are real and detectable only in relation to other objects ... therefore empty space, a void, and so space itself is an unnecessary hypothesis.

That is, Leibnitz introduced a *relational* concept into the Aristotelian world view - what we call *space* is a projection of *relationships* between material bodies into the perceived world whilst what we call *time* is the projection of ordered *change* into the perceived world. Of the three arguments, this latter was the only one to which Clarke had a good objection - essentially that *accelerated motion*, unlike uniform motion, can be perceived *without* reference to external bodies and is therefore, he argued, necessarily perceived with respect to the *absolute space* of Newton. It is of interest to note, however, that in rebutting this particular argument of Leibnitz, Clarke, in the last letter of the correspondence, put his finger directly upon one of the crucial consequences of a relational theory which Leibnitz had apparently not realized (but which Mach much later would) stating as absurd that:

... the parts of a circulating body (suppose the sun) would lose the vis centrifuga arising from their circular motion if all the extrinsic matter around

them were annihilated.

This letter was sent on October 29th 1716 and Leibnitz died on November 14th 1716 so that we were never to know what Leibnitz's response might have been.

Notwithstanding Leibnitz's arguments against the Newtonian conception, nor Berkeley's contemporary criticisms [27], which were very similar to those of Leibnitz and are the direct antecedants of Mach's, the practical success of the Newtonian prescription subdued any serious interest in the matter for the next 150 years or so until Mach himself picked up the torch. In effect, he answered Clarke's response to Leibnitz's second argument by suggesting that the *inertia* of bodies is somehow induced within them by the large-scale distribution of material in the universe:

... I have remained to the present day the only one who insists upon referring the law of inertia to the earth and, in the case of motions of great spatial and temporal extent, to the fixed stars ... [28]

thereby generalizing Leibnitz's conception of a relational universe. Mach was equally clear in expressing his views about the nature of time: in effect, he viewed *time* (specifically Newton's *absolute time*) as a meaningless abstraction. All that we can ever do, he argued in [28], is to measure *change* within one system against *change* in a second system which has been defined as the standard (eg it takes half of one complete rotation of the earth about its own axis to walk thirty miles).

Whilst Mach was clear about the origins of inertia (in the fixed stars), he did not hypothesize any mechanism by which this conviction might be realized and it fell to others to make the attempt - a typical (although incomplete) list might include the names of Einstein [29], Sciama [30], Hoyle & Narlikar and Sachs [31, 32] for approaches based on canonical ideas of spacetime, and the names of Ghosh [33] and Assis [34] for approaches based on quasi-Newtonian ideas.

It is perhaps one of the great ironies of 20thC science that Einstein, having coined the name *Mach's Principle* for Mach's original suggestion and setting out to find a theory which satisfied the newly named Principle, should end up with a theory which, whilst albiet enormously successful, is more an heir to the ideas of Democritus and Newton than to the ideas of Aristotle, Leibnitz and Berkeley. One only has to consider the special case solution of

Minkowski spacetime, which is empty but metrical, to appreciate this fact.

2 Overview

Following in the tradition of Aristotle, Leibnitz, Berkeley and Mach we argue that no consistent cosmology should admit the possibility of an internally consistent discussion of *empty* metrical space & time - unlike, for example, General Relativity which has the empty spacetime of Minkowski as a particular solution. Recognizing that the most simple space & time to visualize is one which is everywhere inertial (which, in any case, approximates the reality of our universe on very large scales) then our worldview is distilled into the elemental question:

Is it possible to conceive a globally inertial space & time which is irreducibly associated with a non-trivial global mass distribution and, if so, what are the properties of this distribution?

In pursuit of this question, this paper falls naturally into three main parts, the first being a development of a Leibnitzian view of physical space, the second being a development of what ‘time’ means in such a physical space and the third being a brief discussion of dynamics in the resulting Leibnitzian universe.

2.1 Leibnitzian physical space: overview

We take the general position of Leibnitz about the relational nature of space to be self-evident and in §3, §4 and §5 develop a quantitative Leibnitzian model of a metric three-space - that is, of a physical three-space whose metric properties are defined entirely in terms of its material content and for which the notion of *empty space* is a meaningless abstraction. This development is notable for the fact that it proceeds from the simple assumption that we can ‘do local physics’ in the conventional way (ie define standard metres and standard seconds and orthogonal reference frames etc for purely local use) and from the primitive and *definitive* knowledge that, in the universe of our experience, the angular sizes of distant objects are (statistically) in a strong inverse relationship with measured cosmological redshifts (a redshift measurement is a local measurement). A purely *deductive* argument - containing no additional hypotheses (beyond the simplifying, but strictly unnecessary, assumption that the large scale universe is statistically unchanging over all

epochs) - then leads to the conclusion that, on very large scales, space necessarily tends to become Euclidean and is supported by a quasi-fractal $D \approx 2$ distribution of universal material.

2.2 Time in a Leibnitzian physical space: overview

The notion of *time* in the Leibnitzian universe is developed in §6, §7, and §8. The assumption that the universe is statistically unchanging over all epochs is equivalent to stating that there is no cosmic time. Therefore, to ‘do’ dynamics in our Leibnitzian universe some concept of local time must be introduced. To keep things simple so that ideas can be properly clarified, we impose the requirement that dynamics means Newtonian dynamics - in essence, that the Third Law must be satisfied. This then leads to a natural definition of time which, within the context of our Leibnitzian universe, can be interpreted to mean that *time* is an explicit measure change in this universe, which very much coincides with Mach’s ideas about the nature of time briefly mentioned in §1.1.

2.3 Dynamics in the Leibnitzian universe: overview

By dynamics here, we mean *gravitational dynamics* and this is discussed in §9 and §10. We show that, on very large scales, all motions tend to become inertial so that, when combined with the result that this large-scale Leibnitzian universe is supported by a quasi-fractal $D \approx 2$ material distribution, the over-all picture is a close reflection of what we actually observe on very large scales.

On small scales in this universe, it becomes clear how Newtonian gravitation emerges in a natural way - although we do not discuss this in detail here. Of equal interest is the fact that for particular limiting values of two parameters, the material in the universe becomes exacty fractal, $D = 2$, and and there is global dynamical equilibrium on all scales. The significance of this is discussed.

3 Leibnitzian physical space

3.1 The qualitative model

In pursuit of the general question raised in the opening paragraph of §2, we shall assume an idealized universe:

- *within which observers can define orthogonal frames locally and make quantitative length measurements/comparisons on local scales only and make measurements of angular diameter and angular position for objects at arbitrary distances;*
- *which, on very large scales, consists of an infinite set of identical discrete galaxies on which there is a primitive property of absolute ordering which allows us to say only that galaxy $\mathcal{G}_0$ is nearer/further than galaxy $\mathcal{G}_1$. There are two obvious consequences of this latter requirement:*
 - *we must be able to see the galaxies and so they must emit light;*
 - *there must be some measurable parameter which allows the judgement of nearer/further. This could be, for example, measured angular diameters or redshifts.*
- *within which there is a primitive temporal ordering property which allows a distinction to be made between before and after. This property will provide a qualitative measure of process or ordered change in the model universe;*
- *within which there is at least one origin about which the distribution of ‘galaxies’ is statistically isotropic - meaning that the results of sampling along arbitrary lines of sight over sufficiently long characteristic ‘times’ are independent of the directions of lines of sight;*
- *within which the distribution of ‘galaxies’ is statistically stationary - meaning that the results of sampling along arbitrary lines of sight over sufficiently long characteristic ‘times’ are independent of sampling epoch.*

3.1.1 Astrophysical spheres and a mass-calibrated metric for radial displacements

We wish to obtain a conception of space & time within which it is *meaningless* to talk about *metric* in the absence of matter. We imagine that we are in the world of our experience and that this world is represented by the primitive model described above. Thus, for the sake of definiteness, we will suppose that the primitive spatial ordering of galaxies can be determined by redshift measurements - in other words, we use only the knowledge that there exists a positive correlation between the (statistical) angular diameters of galaxies and their measured redshifts.

It is then useful to discuss, briefly, the notion of spherical volumes defined on large astrophysical scales in this universe: whilst we can certainly give various precise operational definitions of spherical volumes on local scales, the process of giving such definitions on large scales is decidedly ambiguous. In effect, we have to suppose that redshift measurements are (statistically) isotropic when taken from an arbitrary point within the universe and that they vary monotonically with distance on the large scales we are concerned with. With these assumptions, spherical volumes can be defined (statistically) in terms of redshift measurements - however, their radial calibration in terms of ordinary units (such as metres) becomes increasingly uncertain (and even unknown) on very large redshift scales.

With these ideas in mind, the primary step taken in answer to the elemental question of §3.1 is the recognition that, on large enough scales in the universe of our experience (say $> 30 \text{ Mpc}$), the amount of matter in a given redshift-defined spherical volume in a given epoch can be considered as a well-defined (monotonic) function of the sphere's (redshift) radius.

It follows immediately that a redshift calibration of the radius of an astrophysical sphere has an equivalent mass-defined calibration.

That is, the redshift-radius, R_z say, of any spherical volume, calibrated in terms of mass, can be considered given by:

$$R_z = f(m) \rightarrow \delta R_z \approx f(m + \delta m) - f(m) \quad (1)$$

where m is the mass concerned and f is some well-defined, but unknown, monotonic increasing function of m . Thus, whatever the form of f , we have immediately defined an invariant radial measurement such that it becomes

undefined in the absence of matter - in effect, we have, in principle, a metric which follows Leibnitz in the required sense for any displacement which is *purely radial*.

At this point, we can note that there will be some monotonic relation between a redshift-defined radial displacement R_z and a ‘conventional’ radial measure, R say. Thus, the first of (1) above can be written as

$$R_z \equiv g(R) = f(m) \rightarrow R = g^{-1}(R_z), \quad m = M(R) \leftrightarrow R = M^{-1}(m) \quad (2)$$

where f is a well-defined, but unknown function of m , and g is an arbitrary monotonic increasing function of R , the definition of which provides a calibration of R in terms of R_z .

3.1.2 Definition of a three-dimensional coordinate frame

We now consider how to generalize the foregoing ideas to provide coordinate positions in three-dimensional space for arbitrarily placed distant objects. The method is straightforward: since we have assumed that an observer in the model universe can define a *local* orthogonal frame then he can set up a local rectangular frame with uncalibrated axes, (x^1, x^2, x^3) say, within which he can measure the angular position of arbitrarily placed objects, (θ, ϕ) say. These axes can be supposed extended indefinitely along the lines of sight - but they are completely uncalibrated and cannot be assumed orthogonal on large scales. However, since we have $R = g^{-1}(R_z)$ where R_z is a measured astrophysical quantity, then the observer can provide a global calibration of his (locally) rectangular axes to within a definition of g by *defining*

$$x^1 = R \sin \theta \cos \phi, \quad x^2 = R \sin \theta \sin \phi, \quad x^3 = R \cos \theta. \quad (3)$$

This procedure has the consequence that each of the three axes is calibrated on an astrophysical scale in the same way as R (that is, to within a definition of g) and of guaranteeing that

$$R^2 = (x^1)^2 + (x^2)^2 + (x^3)^2 \rightarrow RdR = x^i dx^j \delta_{ij}.$$

3.1.3 A mass-calibrated metric for arbitrary spatial displacements

We are now in a position to generalize the idea of a mass-calibrated radial displacement to that of a mass-calibration for arbitrary displacements in our

model universe. To this end, we use $m = M(R)$ from (2) and use it in conjunction with (3) to get the mass model:

$$Mass \equiv m = M(R) \equiv M(x^1, x^2, x^3),$$

for our rudimentary universe.

Now consider the normal gradient vector $n_a = \nabla_a M$ and the change in this arising from a displacement dx^k ,

$$dn_a = \nabla_i (\nabla_a M) dx^i, \quad (4)$$

where we assume that the geometrical connections required to give this latter expression an unambiguous meaning are the usual metrical connections - except of course, the metric tensor g_{ab} of our three-space is not yet defined.

Given that $\nabla_a \nabla_b M$ is nonsingular, then (4) provides a 1:1 mapping between the contravariant vector dx^a and the covariant vector dn_a so that, in the absence of any other definition, we can *define* dn_a to be the covariant form of dx^a . In this latter case the metric tensor of our three-space automatically becomes

$$g_{ab} = \nabla_a \nabla_b M \quad (5)$$

which, through the implied metrical connections, is a highly non-linear equation defining g_{ab} to within the specification of the mass model, M . The scalar product $dS^2 \equiv dn_i dx^i \equiv g_{ij} dx^i dx^j$ then provides an invariant measure for the magnitude of the infinitesimal three-space displacement, dx^a .

The units of dS^2 are easily seen to be those of *mass* only and so, in order to make them those of *length*² - as dimensional consistency requires - we define the working invariant as $ds^2 \equiv (2r_0^2/m_0)dS^2$, where r_0 and m_0 are scaling constants for the distance and mass scales respectively and the numerical factor has been introduced for later convenience.

Finally, if we want

$$ds^2 \equiv \left(\frac{r_0^2}{2m_0} \right) dn_i dx^i \equiv \left(\frac{r_0^2}{2m_0} \right) g_{ij} dx^i dx^j \quad (6)$$

to behave sensibly in the sense that $ds^2 > 0$ whenever $|d\mathbf{r}| > 0$ and $ds^2 = 0$ only when $|d\mathbf{r}| = 0$, then we must replace the condition of non-singularity

of g_{ab} by the condition that it is strictly positive definite; in the physical context of the present problem, this will be considered to be a self-evident requirement.

3.2 The connection coefficients

We have assumed that the geometrical connection coefficients can be defined in some sensible way. To do this, we simply note that, in order to define conservation laws (ie to do physics) in a Riemannian space, it is necessary to have a generalized form of Gauss's divergence theorem in the space. This is certainly possible when the connections are defined to be the metrical connections, but it is by no means clear that it is ever possible otherwise. Consequently, the connections are assumed to be metrical and so g_{ab} , given at (5), can be written explicitly as

$$g_{ab} \equiv \nabla_a \nabla_b M \equiv \frac{\partial^2 M}{\partial x^a \partial x^b} - \Gamma_{ab}^k \frac{\partial M}{\partial x^k}, \quad (7)$$

where Γ_{ab}^k are the Christoffel symbols, and given by

$$\Gamma_{ab}^k = \frac{1}{2} g^{kj} \left(\frac{\partial g_{bj}}{\partial x^a} + \frac{\partial g_{ja}}{\partial x^b} - \frac{\partial g_{ab}}{\partial x^j} \right).$$

4 The metric tensor given in terms of the mass model

It is shown, in appendix A, how, for an arbitrarily defined mass model, $m \equiv M(R)$, (7) can be exactly resolved to give an explicit form for g_{ab} in terms of such a general $M(R)$: Defining

$$\mathbf{R} \equiv (x^1, x^2, x^3), \quad \Phi \equiv \frac{1}{2} (\mathbf{R} \cdot \mathbf{R}) = \frac{1}{2} R^2 \quad \text{and} \quad M' \equiv \frac{dM}{d\Phi}$$

then it is found that

$$g_{ab} = A \delta_{ab} + B x^i x^j \delta_{ia} \delta_{jb}, \quad (8)$$

where

$$A \equiv \frac{d_0 M + m_1}{\Phi}, \quad B \equiv -\frac{A}{2\Phi} + \frac{d_0 M' M'}{2A\Phi}.$$

for arbitrary constants d_0 and m_1 where, as inspection of the structure of these expressions for A and B shows, d_0 is a dimensionless scaling factor and m_1 has dimensions of mass. Noting now that M always occurs in the form $d_0M + m_1$, it is convenient to write $\mathcal{M} \equiv d_0M + m_1$, and to write A and B as

$$A \equiv \frac{\mathcal{M}}{\Phi}, \quad B \equiv -\left(\frac{\mathcal{M}}{2\Phi^2} - \frac{\mathcal{M}'\mathcal{M}'}{2d_0\mathcal{M}}\right). \quad (9)$$

5 An invariant calibration of the radial scale

So far, we have assumed an arbitrary calibration for the radial scale; that is, since, by (2), $R_z = f(m)$ for unknown f and since $R = g^{-1}(R_z)$ for arbitrary monotonic g , then $R = g^{-1}\{f(m)\}$ is uncalibrated. In the following, we show that the *geodesic radial scale* provides a unique calibration for R in terms of m and leads directly to an understanding of fractality, $D \approx 2$, in our observed universe.

5.1 The geodesic radial scale

Using (8) and (9) in (6), and applying the identities $x^i dx^j \delta_{ij} \equiv R dR$ and $\Phi \equiv R^2/2$, we find, for an arbitrary displacement $d\mathbf{x}$, the invariant measure:

$$ds^2 = \left(\frac{R_0^2}{2m_0}\right) \left\{ \frac{\mathcal{M}}{\Phi} dx^i dx^j \delta_{ij} - \Phi \left(\frac{\mathcal{M}}{\Phi^2} - \frac{\mathcal{M}'\mathcal{M}'}{d_0\mathcal{M}} \right) dR^2 \right\},$$

which is valid for the arbitrary calibration $R = g^{-1}\{f(m)\}$. If the displacement $d\mathbf{x}$ is now constrained to be purely radial, then we find

$$ds^2 = \left(\frac{R_0^2}{2m_0}\right) \left\{ \Phi \left(\frac{\mathcal{M}'\mathcal{M}'}{d_0\mathcal{M}} \right) dR^2 \right\}.$$

Use of $\mathcal{M}' \equiv d\mathcal{M}/d\Phi$ and $\Phi \equiv R^2/2$ reduces this latter relationship to

$$\begin{aligned} ds^2 &= \frac{R_0^2}{d_0 m_0} (d\sqrt{\mathcal{M}})^2 \rightarrow ds = \frac{R_0}{\sqrt{d_0 m_0}} d\sqrt{\mathcal{M}} \rightarrow \\ s &= \frac{R_0}{\sqrt{d_0 m_0}} \left(\sqrt{\mathcal{M}} - \sqrt{\mathcal{M}_0} \right), \quad \text{where } \mathcal{M}_0 \equiv \mathcal{M}(s=0) \end{aligned}$$

which defines the invariant magnitude of an arbitrary *radial* displacement from the origin purely in terms of the mass-model representation $\mathcal{M} \equiv$

$d_0M + m_1$. By definition, this s is the invariant measure for an arbitrary finite radial displacement - in other words, it is the natural measure of radial displacement. Thus, by setting $R \equiv s$, we identify R with the natural measure of radial displacement, and thereby provide it with a unique calibration.

To summarize, the natural mass-defined calibration for the radial scale is given by

$$R = \frac{R_0}{\sqrt{d_0 m_0}} \left(\sqrt{\mathcal{M}} - \sqrt{\mathcal{M}_0} \right), \quad (10)$$

where $\mathcal{M}_0$ is the value of $\mathcal{M}$ at $R = 0$.

5.2 The Euclidean metric and a fractal $D = 2$ mass distribution

Using $\mathcal{M} \equiv d_0M + m_1$ and noting that $M(R = 0) = 0$ necessarily, then $\mathcal{M}_0 = m_1$ and so (10) can be equivalently arranged as

$$\mathcal{M} = \left[\frac{\sqrt{d_0 m_0}}{R_0} R + \sqrt{m_1} \right]^2. \quad (11)$$

Using $\mathcal{M} \equiv d_0M + m_1$ again, then the mass-distribution function can be expressed in terms of the invariant radial displacement as

$$M = m_0 \left(\frac{R}{R_0} \right)^2 + 2\sqrt{\frac{m_0 m_1}{d_0}} \left(\frac{R}{R_0} \right) \quad (12)$$

which, for the particular case $m_1 = 0$ becomes $M = m_0(R/R_0)^2$. Reference to (8) shows that, with this mass distribution and $d_0 = 1$, then $g_{ab} = \delta_{ab}$ so that the three-space space becomes ordinary Euclidean space. Thus, whilst we have yet to show that a globally inertial space can be associated with a non-trivial global matter distribution (since no temporal dimension, and hence no dynamics, has been introduced), we have shown that a globally *Euclidean* space can be associated with a non-trivial matter distribution, and that this distribution is necessarily fractal with $D = 2$.

Note also that, on a large enough scale and for *arbitrary* values of m_1 , (12) shows that radial distance varies as the square-root of mass from the chosen origin - or, equivalently, the mass varies as R^2 . Consequently, on sufficiently

large scales Euclidean space is irreducibly related to a quasi-fractal, $D = 2$, matter distributions. Since $M/R^2 \approx m_0/R_0^2$ on a large enough scale then, for the remainder of this paper, the notation $g_0 \equiv m_0/R_0^2$ is employed.

6 The temporal dimension

So far, the concept of ‘time’ has only entered the discussion in a qualitative way in §1.1 - it has not entered in any quantitative way and, until it does, there can be no discussion of dynamical processes.

Since, in its most general definition, time is a parameter which orders change within a system, then a necessary pre-requisite for its quantitative definition is a notion of change within the universe. The most simple notion of change which can be defined in the universe is that of changing relative spatial displacements of the objects within it. Since our model universe is populated solely by identical primitive ‘galaxies’ then, in effect, all change is gravitational change. This fact is incorporated into the cosmology to be derived by constraining all particle displacements to satisfy the Weak Equivalence Principle. We are then led to a Lagrangian description of particle motions in which the Lagrange density is degree zero in its temporal-ordering parameter. From this, it follows that the corresponding Euler-Lagrange equations form an *incomplete* set.

The origin of this problem traces back to the fact that, because the Lagrangian density is degree zero in the temporal ordering parameter, it is then invariant with respect to any transformation of this parameter which preserves the ordering. This implies that, in general, temporal ordering parameters cannot be identified directly with physical time - they merely share one essential characteristic. This situation is identical to that encountered in the Lagrangian formulation of General Relativity; there, the situation is resolved by defining the concept of ‘particle proper time’. In the present case, this is not an option because the notion of particle proper time involves the prior definition of a system of observer’s clocks - so that some notion of clock-time is factored into the prior assumptions upon which General Relativity is built.

In the present case, it turns out that the isotropies already imposed on the

system conspire to provide an automatic resolution of the problem which is consistent with the already assumed interpretation of ‘time’ as a measure of ordered change in the model universe. To be specific, it turns out that the elapsed time associated with any given particle displacement is proportional, via a scalar field, to the invariant spatial measure attached to that displacement. Thus, physical time is defined directly in terms of the invariant measures of *process* within the model universe and, furthermore, local conditions affect clock rates.

7 Equations of motion for generalized time parameter

Firstly, and as already noted, we are assuming that, within our model universe, all motions are gravitational, and we model this circumstance by constraining all such motions to satisfy the Weak Equivalence Principle by which we mean that the trajectory of a body is independent of its internal constitution. This constraint can be expressed as:

C1 Particle trajectories are independent of the specific mass values of the particles concerned;

Secondly, given the isotropy conditions imposed on the model universe from the chosen origin, symmetry arguments lead to the conclusion that the net action of the whole universe of particles acting on any given single particle is such that any net acceleration of the particle must always appear to be directed through the coordinate origin. Note that this conclusion is *independent* of any notions of retarded or instantaneous action. This constraint can then be stated as:

C2 Any acceleration of any given material particle must necessarily be along the line connecting the particular particle to the coordinate origin.

Now suppose p and q are two arbitrarily chosen point coordinates on the trajectory of the chosen particle, and suppose that (6) is integrated between these points to give the scalar invariant

$$I(p, q) = \int_p^q \left(\frac{1}{\sqrt{2g_0}} \right) \sqrt{dn_i dx^i} \equiv \int_p^q \left(\frac{1}{\sqrt{2g_0}} \right) \sqrt{g_{ij} dx^i dx^j}. \quad (13)$$

Then, in accordance with the foregoing interpretation, $I(p, q)$ gives a scalar record of how the particle has moved between p and q defined with respect to

the particle's continually changing relationship with the mass model, $M(R)$.

If $I(p, q)$ is now minimized with respect to choice of the trajectory connecting p and q , then this minimizing trajectory can be interpreted as a geodesic in the Riemannian space which has g_{ab} as its metric tensor. Given that g_{ab} is defined in terms of the mass model $M(R)$ - the existence of which is independent of any notion of 'inertial mass', then the existence of the metric space, and of geodesic curves within it, is likewise explicitly independent of any concept of inertial-mass. It follows that the identification of the particle trajectory $\mathbf{r}$ with these geodesics means that particle trajectories are similarly independent of any concept of inertial mass, and can be considered as the modelling step defining that general subclass of trajectories which conform to that characteristic phenomenology of gravitation defined by condition **C1** above.

The geodesic equations in the space with the metric tensor (8) can now be obtained, in the usual way, by defining the Lagrangian density

$$\mathcal{L} \equiv \left(\frac{1}{\sqrt{2g_0}} \right) \sqrt{g_{ij} \dot{x}^i \dot{x}^j} = \left(\frac{1}{\sqrt{2g_0}} \right) \left(A (\dot{\mathbf{R}} \cdot \dot{\mathbf{R}}) + B \dot{\Phi}^2 \right)^{1/2}, \quad (14)$$

where $\dot{x}^i \equiv dx^i/dt$, etc., and writing down the Euler-Lagrange equations

$$\begin{aligned} 2A\ddot{\mathbf{R}} + \left(2A'\dot{\Phi} - 2\frac{\dot{\mathcal{L}}}{\mathcal{L}}A \right) \dot{\mathbf{R}} + \left(B'\dot{\Phi}^2 + 2B\ddot{\Phi} - A'(\dot{\mathbf{R}} \cdot \dot{\mathbf{R}}) - 2\frac{\dot{\mathcal{L}}}{\mathcal{L}}B\dot{\Phi} \right) \mathbf{R} \\ = 0, \end{aligned} \quad (15)$$

where $\dot{\mathbf{R}} \equiv d\mathbf{R}/dt$ and $A' \equiv dA/d\Phi$, etc. By identifying particle trajectories with geodesic curves, this equation is now interpreted as the equation of motion, referred to the chosen origin, of a single particle satisfying condition **C1** above.

However, noting that the variational principle, (13), is of order zero in its temporal ordering parameter, we can conclude that the principle is invariant with respect to arbitrary transformations of this parameter; in turn, this means that the temporal ordering parameter cannot be identified with physical time. This problem manifests itself formally in the statement that the equations of motion (15) do not form a complete set, so that it becomes necessary to specify some extra condition to close the system.

A similar circumstance arises in General Relativity when the equations of motion are derived from an action integral which is formally identical to (13). In that case, the system is closed by specifying the arbitrary time parameter to be the ‘proper time’, so that

$$d\tau = \mathcal{L}(x^j, dx^j) \rightarrow \mathcal{L}(x^j, \frac{dx^j}{d\tau}) = 1, \quad (16)$$

which is then considered as the necessary extra condition required to close the system. In the present circumstance, we are rescued by the, as yet, unused condition **C2**.

8 Physical time

8.1 Completion of equations of motion

Consider **C2**, which states that any particle accelerations must necessarily be directed through the coordinate origin. This latter condition simply means that the equations of motion must have the general structure

$$\ddot{\mathbf{R}} = G(t, \mathbf{R}, \dot{\mathbf{R}})\mathbf{R},$$

for scalar function $G(t, \mathbf{R}, \dot{\mathbf{R}})$. In other words, (15) satisfies condition **C2** if the coefficient of $\dot{\mathbf{R}}$ is zero, so that

$$\left(2A'\dot{\Phi} - 2\frac{\dot{\mathcal{L}}}{\mathcal{L}}A\right) = 0 \rightarrow \frac{A'}{A}\dot{\Phi} = \frac{\dot{\mathcal{L}}}{\mathcal{L}} \rightarrow \mathcal{L} = k_0 A, \quad (17)$$

for arbitrary constant k_0 which is necessarily positive since $A > 0$ and $\mathcal{L} > 0$. The condition (17), which guarantees (**C2**), can be considered as the condition required to close the incomplete set (15) and is directly analogous to (16), the condition which defines ‘proper time’ in General Relativity.

8.2 Physical time defined as process

Equation (17) can be considered as that equation which removes the pre-existing arbitrariness in the ‘time’ parameter by *defining* physical time:- from

(17) and (14) we have

$$\begin{aligned}\mathcal{L}^2 &= k_0^2 A^2 \rightarrow A(\dot{\mathbf{R}} \cdot \dot{\mathbf{R}}) + B\dot{\Phi}^2 = 2g_0 k_0^2 A^2 \rightarrow \\ g_{ij}\dot{x}^i\dot{x}^j &= 2g_0 k_0^2 A^2\end{aligned}\tag{18}$$

so that, in explicit terms, physical time is *defined* by the relation

$$dt^2 = \left(\frac{1}{2g_0 k_0^2 A^2} \right) g_{ij} dx^i dx^j, \quad \text{where } A \equiv \frac{\mathcal{M}}{\Phi}.\tag{19}$$

In short, the elapsing of time is given a direct physical interpretation in terms of the process of *displacement* in the model universe.

Finally, noting that, by (19), the dimensions of k_0^2 are those of $L^6/[T^2 \times M^2]$, then the fact that $g_0 \equiv m_0/R_0^2$ (cf §5) suggests the change of notation $k_0^2 \propto v_0^2/g_0^2$, where v_0 is a constant having the dimensions (but not the interpretation) of ‘velocity’. So, as a means of making the dimensions which appear in the development more transparent, it is found convenient to use the particular replacement $k_0^2 \equiv v_0^2/(4d_0^2 g_0^2)$. With this replacement, the *definition* of physical time, given at (19), becomes

$$dt^2 = \left(\frac{4d_0^2 g_0}{v_0^2 A^2} \right) g_{ij} dx^i dx^j.\tag{20}$$

Since, as is easily seen from the definition of g_{ab} given in §4, $g_{ij} dx^i dx^j$ is necessarily finite and non-zero for a non-trivial displacement $d\mathbf{R}$

8.3 The necessity of $v_0^2 \neq 0$

Equation (20) provides a definition of physical time in terms of basic process (displacement) in the model universe. Since the parameter v_0^2 occurs nowhere else, except in its explicit position in (20), then it is clear that setting $v_0^2 = 0$ is equivalent to physical time becoming undefined. Therefore, of necessity, $v_0^2 \neq 0$ and all non-zero finite displacements are associated with a non-zero finite elapsed physical time.

9 The cosmological potential

The model is most conveniently interpreted when expressed in potential terms and so, in the following, it is shown how this is done.

9.1 The equations of motion: potential form

When (17) is used in (15) we get:

$$2A\ddot{\mathbf{R}} + \left(B'\dot{\Phi}^2 + 2B\ddot{\Phi} - A'(\dot{\mathbf{R}} \cdot \dot{\mathbf{R}}) - 2\frac{A'}{A}B\dot{\Phi}^2 \right) \mathbf{R} = 0. \quad (21)$$

Suppose we define a function V according to $V \equiv C_0 - (\dot{\mathbf{R}} \cdot \dot{\mathbf{R}})/2$, for some arbitrary constant C_0 ; then, by (18)

$$V \equiv C_0 - \frac{1}{2}(\dot{\mathbf{R}} \cdot \dot{\mathbf{R}}) = C_0 - \frac{v_0^2}{4d_0^2g_0}A + \frac{B}{2A}\dot{\Phi}^2, \quad (22)$$

where A and B are defined at (9). With unit vector, $\hat{\mathbf{R}}$, then appendix B shows how this function can be used to express (21) in the potential form

$$\ddot{\mathbf{R}} = -\frac{dV}{dr}\hat{\mathbf{R}} \quad (23)$$

so that V is a potential function, and C_0 is the arbitrary constant usually associated with a potential function.

9.2 The potential function, V , as a function of R

From (22), we have

$$2C_0 - 2V = \dot{R}^2 + R^2\dot{\theta}^2 = \frac{v_0^2}{2d_0^2g_0}A - \frac{B}{A}R^2\dot{R}^2$$

so that V is effectively given in terms of R and $\dot{R}$. In order to clarify things further, we now eliminate the explicit appearance of $\dot{R}$. Since all forces are central, then angular momentum is conserved; consequently, after using conserved angular momentum, h , and the definitions of A , B and $\mathcal{M}$ given in §4, the foregoing equations can be written as

$$\begin{aligned} 2C_0 - 2V = \\ \dot{R}^2 + R^2\dot{\theta}^2 = v_0^2 + \frac{4v_0^2}{R}\sqrt{\frac{m_1}{d_0g_0}} + \frac{d_0 - 1}{R^2}\left(\frac{6m_1v_0^2}{d_0^2g_0} - h^2\right) \\ + \frac{2}{R^3}\sqrt{\frac{d_0m_1}{g_0}}\left(\frac{2m_1v_0^2}{d_0^2g_0} - h^2\right) + \frac{1}{R^4}\frac{m_1}{g_0}\left(\frac{m_1v_0^2}{d_0^2g_0} - h^2\right) \end{aligned} \quad (24)$$

so that $V(R)$ is effectively given by the right-hand side of (24). Various interesting features of this potential function, which go beyond our immediate interest, are discussed in appendix C.

10 The fractal $D = 2$ inertial universe

The model we have developed allows either a perfectly fractal $D = 2$ inertial universe as a specific limiting case, or a quasi-fractal $D \approx 2$, almost inertial universe on sufficiently large scales. In either case, the notion of an empty metrical universe is meaningless, so that the question originally posed in §2 is finally answered. We consider each case in turn.

10.1 The perfectly fractal $D = 2$ universe

Reference to (24) shows that the parameter choice $m_1 = 0$ and $d_0 = 1$ makes the potential function constant everywhere so that there is a global dynamical equilibrium; similarly, (12) shows how this equilibrium universe is supported by a mass distribution which is necessarily distributed as an exact fractal with $D = 2$.

A more detailed analysis is illuminating: specifically, for the case $m_1 = 0$, $d_0 = 1$, the definition M at (12) together with the definitions of A and B in §4 give

$$A = \frac{2m_0}{R_0^2}, \quad B = 0$$

so that, by (22) (remembering that $g_0 \equiv m_0/R_0^2$) we have

$$(\dot{\mathbf{R}} \cdot \dot{\mathbf{R}}) = v_0^2 \tag{25}$$

for all displacements in the model universe. It is (almost) natural to assume that the constant v_0^2 in (25) simply refers to the constant velocity of any given particle, and likewise to assume that this can differ between particles. However, each of these assumptions would be wrong since - as we now show - v_0^2 is, firstly, more properly interpreted as a conversion factor from spatial to temporal units and, secondly, is a *global* constant which applies equally to all particles.

To understand these points, we begin by noting that (25) is a special case of (18) and so, by (19), is more accurately written as

$$dt^2 = \frac{1}{v_0^2} (d\mathbf{R} \cdot d\mathbf{R}) \tag{26}$$

which, by the considerations of §8.2, we recognize as the *definition* of the elapsed time experienced by any particle undergoing a spatial displacement $d\mathbf{r}$ in the model inertial universe. Since this universe is isotropic about all points, then there is nothing which can distinguish between two separated particles (other than their separateness) undergoing displacements of equal magnitudes; consequently, each must be considered to have experienced equal elapsed times. It follows from this that v_0^2 is not to be considered as a locally defined particle velocity, but is a *globally* defined constant which has the effect of converting between spatial and temporal units of measurement.

We now see that the model inertial universe, with (26) as a global relationship, bears a close formal resemblance to a universe filled purely with Einsteinien photons - the difference being, of course, that the particles in the model inertial universe are assumed to be countable and to have mass properties. This formal resemblance means that the idealized inertial universe can be likened to a quasi-photon fractal, $D = 2$, gas universe.

10.2 The quasi-fractal $D \approx 2$ universe

On large scales, and for arbitrary choice of the parameters, (12) shows that $M \sim R^2$ on sufficiently large scales whilst (24) shows that accelerations $\rightarrow 0$ on these scales. Using arguments similar to those used above, we can also show that the elapsed time experienced by particles undergoing displacements in this universe becomes a function of the displacement itself and the local environment.

10.3 Implications for theories of gravitation

Given that gravitational phenomena are usually considered to arise as mass-driven perturbations of flat inertial backgrounds, then the foregoing results - to the effect that the inertial background is necessarily associated with a non-trivial fractal matter distribution - must necessarily give rise to completely new perspectives about the nature and properties of gravitational phenomena. However, such considerations take us beyond the immediate concerns of this paper.

11 Summary and Conclusions

In practice, and insofar as we can determine, the universe of our experience is asymptotically inertial on the very large scales at which quasi-fractal $D \approx 2$ behaviour appears to dominate. We adopted the tentative hypothesis that this quasi-fractal behaviour will be found to extend to all observable (large) scales and that it is a persistent signature reflecting the fundamental nature of the universe as a place in which the ideas Aristotle, Leibnitz, Berkeley and Mach are a fair reflection of the reality. If this is the case, then it should be possible to produce a theoretical description of such a universe so that we were led to ask the question:

Is it possible to associate a globally inertial space & time with a non-trivial global matter distribution and, if it is, what are the fundamental properties of this distribution?

We analysed the question in the context of a simple model universe which was assumed to possess only the primitive concepts of *nearer/further* and of *before/after*. We then showed how quantitative ideas of spatial and temporal metrics could be defined entirely in terms of the universal material and that, as a direct consequence:

- on sufficiently large scales in this universe and for an arbitrary choice of the parameters, the distribution of material becomes quasi-fractal with $D \approx 2$ (cf equation (11) whilst accelerations tend to zero (inertial behaviour on large scales, cf (24));
- In an ideal limiting case of two particular parameters taking certain exact values, a globally inertial space & time is irreducibly related to a fractal, $D = 2$, distribution of material within the model universe.

However, the latter ideal inertial universe is distinguished in the sense that whilst all the particles within it have arbitrarily directed motions, the particle velocities all have *equal* magnitude. In this sense, this limiting case globally inertial model universe is more accurately to be considered as a quasi-photon gas universe than the universe of our macroscopic experience. In other words, it looks more like a crude model of a material vacuum than the universe of our direct experience.

A A Resolution of the Metric Tensor

The general system is given by

$$g_{ab} = \frac{\partial^2 M}{\partial x^a \partial x^b} - \Gamma_{ab}^k \frac{\partial M}{\partial x^k},$$

$$\Gamma_{ab}^k \equiv \frac{1}{2} g^{kj} \left(\frac{\partial g_{bj}}{\partial x^a} + \frac{\partial g_{ja}}{\partial x^b} - \frac{\partial g_{ab}}{\partial x^j} \right),$$

and the first major problem is to express g_{ab} in terms of the reference scalar, M . The key to this is to note the relationship

$$\frac{\partial^2 M}{\partial x^a \partial x^b} = M' \delta_{ab} + M'' x^a x^b,$$

where $M' \equiv dM/d\Phi$, $M'' \equiv d^2M/d\Phi^2$ and $\Phi \equiv R^2/2$, since this immediately suggests the general structure

$$g_{ab} = A \delta_{ab} + B x^a x^b, \quad (27)$$

for unknown functions, A and B . It is easily found that

$$g^{ab} = \frac{1}{A} \left[\delta_{ab} - \left(\frac{B}{A + 2B\Phi} \right) x^a x^b \right]$$

so that, with some effort,

$$\Gamma_{ab}^k = \frac{1}{2A} H_1 - \left(\frac{B}{2A(A + 2B\Phi)} \right) H_2$$

where

$$H_1 = A'(x^a \delta_{bk} + x^b \delta_{ak} - x^k \delta_{ab})$$

$$+ B' x^a x^b x^k + 2B \delta_{ab} x^k$$

and

$$H_2 = A'(2x^a x^b x^k - 2\Phi x^k \delta_{ab})$$

$$+ 2\Phi B' x^a x^b x^k + 4\Phi B x^k \delta_{ab}.$$

Consequently,

$$g_{ab} = \frac{\partial^2 M}{\partial x^a \partial x^b} - \Gamma_{ab}^k \frac{\partial M}{\partial x^k} \equiv \delta_{ab} M' \left(\frac{A + A'\Phi}{A + 2B\Phi} \right) + x^a x^b \left(M'' - M' \left(\frac{A' + B'\Phi}{A + 2B\Phi} \right) \right).$$

Comparison with (27) now leads directly to

$$\begin{aligned} A &= M' \left(\frac{A + A'\Phi}{A + 2B\Phi} \right) = M' \left(\frac{(A\Phi)'}{A + 2B\Phi} \right), \\ B &= M'' - M' \left(\frac{A' + B'\Phi}{A + 2B\Phi} \right). \end{aligned}$$

The first of these can be rearranged as

$$B = \frac{M'}{2\Phi} \left(\frac{(A\Phi)'}{A} \right) - \frac{A}{2\Phi}$$

or as

$$\left(\frac{M'}{A + 2B\Phi} \right) = \frac{A}{(A\Phi)'},$$

and these expressions can be used to eliminate B in the second equation. After some minor rearrangement, the resulting equation is easily integrated to give, finally,

$$A \equiv \frac{d_0 M + m_1}{\Phi}, \quad B \equiv -\frac{A}{2\Phi} + \frac{d_0 M' M'}{2A\Phi}.$$

B Conservative Form of Equations of Motion

From (22), we have

$$V \equiv -\frac{1}{2} (\dot{\mathbf{R}} \cdot \dot{\mathbf{R}}) = -\frac{k_0^2 A}{2} + \frac{B}{2A} \dot{\Phi}^2, \quad (28)$$

from which we easily find

$$\frac{dV}{dR} \equiv \frac{\partial V}{\partial R} + \frac{\partial V}{\partial \dot{R}} \frac{\ddot{R}}{\dot{R}}$$

$$= \frac{-k_0^2 A'}{2} R + \frac{\dot{\Phi}^2 R}{2A} \left(B' - \frac{A'B}{A} \right) + \frac{B}{A} (R\dot{R}^2 + R^2\ddot{R})$$

where we remember that $A' \equiv dA/d\Phi$ etc. Since $\dot{R}^2 + R\ddot{R} = \ddot{\Phi}$, then the above leads to

$$\frac{dV}{dR} \hat{\mathbf{R}} = \left(\frac{-k_0^2 A'}{2} + \frac{B'}{2A} \dot{\Phi}^2 - \frac{A'B}{2A^2} \dot{\Phi}^2 + \frac{B}{A} \ddot{\Phi} \right) \mathbf{R}.$$

Writing (23) as

$$2A\ddot{\mathbf{R}} + 2A\frac{dV}{dR}\hat{\mathbf{R}} = 0,$$

and using the above expression, we get the equation of motion as

$$2A\ddot{\mathbf{R}} + \left(-k_0^2 AA' + B'\dot{\Phi}^2 - \frac{A'B}{A}\dot{\Phi}^2 + 2B\ddot{\Phi} \right) \mathbf{R} = 0. \quad (29)$$

Finally, from (28), we have

$$k_0^2 A = \frac{B}{A} \dot{\Phi}^2 + (\dot{\mathbf{R}} \cdot \dot{\mathbf{R}}),$$

which, when substituted into (29), gives (21).

C Outline analysis of the potential function

It is quite plain from (24) that, for any $m_1 \neq 0$, then the model universe has a preferred centre and that the parameter m_1 (which has dimensions of mass) plays a role in the potential V which is analogous to the source mass in a Newtonian spherical potential - that is, the parameter m_1 can be identified as the mass of the potential source in the model universe. However, setting $m_1 = 0$ is not sufficient to guarantee a constant potential field since any $d_0 \neq 1$ also provides the model universe with a preferred centre. The role of d_0 is most simply discussed in the limiting case of $m_1 = 0$: in this case, the second equation of (24) becomes

$$\dot{R}^2 + R^2\dot{\theta}^2 = v_0^2 - (d_0 - 1)\frac{h^2}{R^2}. \quad (30)$$

If $d_0 < 1$ then $|\dot{\mathbf{R}}| \rightarrow \infty$ as $R \rightarrow 0$ so that a singularity exists. Conversely, remembering that $v_0^2 > 0$ (cf §8.3) then, if $d_0 > 1$, equation (30) restricts real

events to the exterior of the sphere defined by $R^2 = (d_0 - 1)h^2/v_0^2$. In this case, the singularity is avoided and the central ‘massless particle’ is given the physical property of ‘finite extension’. In the more realistic case for which $m_1 > 0$, reference to (24) shows that the $R = 0$ singularity is completely avoided whenever $h^2 > m_1 v_0^2/d_0^2 g_0$ since then a ‘finite extension’ property for the central massive particle always exists. Conversely, a singularity will necessarily exist whenever $h^2 \leq m_1 v_0^2/d_0^2 g_0$.

In other words, the model universe has a preferred centre when either $m_1 > 0$, in which case the source of the potential is a massive central particle having various properties depending on the value of d_0 , or when $m_1 = 0$ and $d_1 \neq 0$.

References

- [1] Charlier, C.V.L., 1908, *Astronomi och Fysik* 4, 1 (Stockholm)
- [2] Charlier, C.V.L., 1922, *Ark. Mat. Astron. Physik* 16, 1 (Stockholm)
- [3] Charlier, C.V.L., 1924, *Proc. Astron. Soc. Pac.* 37, 177
- [4] De Vaucouleurs, G. 1970 *Science* 167, 1203
- [5] Peebles, P.J.E. 1980 *The Large Scale Structure of the Universe*, Princeton University Press, Princeton, NJ.
- [6] Baryshev, Yu V., Sylos Labini, F., Montuori, M., Pietronero, L. 1995 *Vistas in Astronomy* 38, 419
- [7] Huchra, J., Davis, M., Latham, D., Tonry, J. 1983 *ApJS* 52, 89
- [8] Giovanelli, R., Haynes, M.P., Chincarini, G.L. 1986 *ApJ* 300, 77
- [9] De Lapparent, V., Geller, M.J., Huchra, J.P. 1988 *ApJ* 332, 44
- [10] Broadhurst, T.J., Ellis, R.S., Koo, D.C., Szalay, A.S. 1990 *Nature* 343, 726
- [11] Da Costa, L.N., Geller, M.J., Pellegrini, P.S., Latham, D.W., Fairall, A.P., Marzke, R.O., Willmer, C.N.A., Huchra, J.P., Calderon, J.H., Ramella, M., Kurtz, M.J. 1994 *ApJ* 424, L1

- [12] Vettolani, G., et al., in: *Proc. of Schloss Rindberg Workshop: Studying the Universe With Clusters of Galaxies*
- [13] Joyce, M., Montuori, M., Labini, F.S. 1999 *ApJ* 514, L5
- [14] Labini, F.S., Montuori, M. 1998 *A&A* 331, 809
- [15] Labini, F.S., Montuori, M., Pietronero, L. 1998 *Phys. Lett.* 293, 62
- [16] Wu, K.K.S., Lahav, O., Rees, M.J., 1999, *Nature* 397, 225
- [17] Scaramella, R., Guzzo, L., Zamorani, G., Zucca, E., Balkowski, C., Blanchard, A., Cappi, A., Cayatte, V., Chincarini, G., Collins, C., Fiorani, A., Maccagni, D., MacGillivray, H., Maurogordato, S., Merighi, R., Mignoli, M., Proust, D., Ramella, M., Stirpe, G.M., Vettolani, G. 1998 *A&A* 334, 404
- [18] Martinez, V.J., PonsBorderia, M.J., Moyeed, R.A., Graham, M.J. 1998 *MNRAS* 298, 1212
- [19] Joyce, M., Montuori, M., Labini, F.S., Pietronero, L. 1999 *A&A* 344, 387
- [20] Labini, F.S., Gabrielli, A., 2000, *Scaling and fluctuations in galaxy distribution: two tests to probe large scale structures*, astro-ph0008047
- [21] Gabrielli, A., Sylos Labini, F., 2001, *Europhys. Lett.* 54 (3), 286
- [22] Newton, I., *Principia: Mathematical Principles of Natural Philosophy*, translated by I.B. Cohen & A. Whitman, University of California Press, 1999
- [23] Aristotle, *Categories*, in R. McKeon, *The Basic Works of Aristotle*, Random House, New York, 1941.
- [24] Aristotle, *Physics*, translated by P.H. Wicksteed & F.M. Cornford, Leob Classical Library, Harvard University Press, Cambridge, 1929
- [25] Leibnitz, G.W., *Philosophical Essays*, Edited and translated by R. Ariew & D.Garber, Hackett Publishing Co, Indianapolis, 1989
- [26] Alexander, H.G., *The Leibnitz-Clarke Correspondence*, Manchester University Press, 1984.

- [27] Berkeley, G., *Of motion, or the principle and nature of motion and the cause of the communication of motions* (Latin: *De Motu*) in M.R.Ayers (ed) *George Berkeley's Philosophical Works*, Everyman, London, 1992.
- [28] Mach, E., *The Science of Mechanics - a Critical and Historical Account of its Development* Open Court, La Salle, 1960
- [29] Einstein, A., *The Foundations of the General Theory of Relativity*, in A. Einstein, H.A.Lorentz, H.Weyl & H.Minkowski, *The Principle of Relativity*, Dover, New York, 1952
- [30] Sciama, D.W., 1953, *On the Origin of Inertia*, Mon. Not. Roy. Astron. Soc, 113, 34
- [31] Sachs, M., *General Relativity and Matter*, Reidal, Dordrecht, 1982
- [32] Sachs, M., *Quantum Mechanics from General Relativity*, Reidal, Dordrecht 1986
- [33] Ghosh, A., *Origin of Inertia*, Apeiron, Montreal, 2000
- [34] Assis, A.K.T., *Relational Mechanics*, Apeiron, Montreal, 1999